

University of Groningen

**Nucleotide sequences of the genes encoding fructosebisphosphatase and phosphoribulokinase from *Xanthobacter flavus* H4-14**

Meijer, Wilhelmus; Enequist, H.G.; Terpstra, Peter; Dijkhuizen, L.

*Published in:*  
Journal of general microbiology

*DOI:*  
[10.1099/00221287-136-11-2225](https://doi.org/10.1099/00221287-136-11-2225)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1990

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Meijer, W., Enequist, H. G., Terpstra, P., & Dijkhuizen, L. (1990). Nucleotide sequences of the genes encoding fructosebisphosphatase and phosphoribulokinase from *Xanthobacter flavus* H4-14. *Journal of general microbiology*, 136(11), 2225-2230. <https://doi.org/10.1099/00221287-136-11-2225>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## Nucleotide sequences of the genes encoding fructosebisphosphatase and phosphoribulokinase from *Xanthobacter flavus* H4-14

W. G. MEIJER,<sup>1</sup>† H. G. ENEQUIST,<sup>2</sup> P. TERPSTRA<sup>2</sup> and L. DIJKHUIZEN<sup>1</sup>\*

<sup>1</sup>Department of Microbiology, University of Groningen, Kerklaan 30, 9751 NN Haren, The Netherlands

<sup>2</sup>Department of Biochemistry, University of Groningen, Nijenborgh 16, 9747 AG Groningen, The Netherlands

(Received 6 July 1990; accepted 9 August 1990)

---

The genes encoding fructosebisphosphatase and phosphoribulokinase present on a 2.5 kb *Sal*I fragment from *Xanthobacter flavus* H4-14 were sequenced. Two large open reading frames (ORFs) were identified, preceded by plausible ribosome-binding sites. The ORFs were transcribed in the same direction and were separated by 39 base pairs. They encoded proteins of 364 and 291 amino acids, with molecular masses of 38739 and 33409 Da, respectively. The ORFs were identified as the genes encoding FBPase and PRK, respectively, on the basis of similarity with FBPase and PRK sequences from other sources.

---

### Introduction

Many organisms, including higher plants, grow autotrophically using the Calvin cycle for CO<sub>2</sub> fixation. Three enzymic steps are considered to be unique to this pathway. Phosphoribulokinase (PRK) catalyses the phosphorylation of ribulose 5-phosphate (RuMP) by ATP, yielding ribulose 1,5-bisphosphate (RuBP) and ADP. The RuBP formed is carboxylated by ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisC/O), leading to the formation of two molecules of 3-phosphoglycerate. RuMP is regenerated by a series of enzymes which, with the exception of sedoheptulosebisphosphatase, also function in other pathways, e.g. during gluconeogenesis. The latter enzyme is bifunctional, also displaying fructosebisphosphatase (FBPase) activity (Tabita, 1988).

The genes encoding FBPase (*cfxF*) and PRK (*cfxP*) have been cloned from the autotrophic bacteria *Alcaligenes eutrophus*, *Rhodobacter sphaeroides* and *Xanthobacter flavus* H4-14. In these organisms the *cfxF* and *cfxP*

genes are cotranscribed in the order *cfxF*–*cfxP* (Gibson & Tabita, 1987, 1988; Hallenbeck & Kaplan, 1987; Klintworth *et al.*, 1988; Meijer *et al.*, 1990*b*). The nucleotide sequences of a number of FBPase-encoding genes have been reported. However, the only Calvin cycle *cfxF* gene that has been sequenced is the one from wheat chloroplast (Raines *et al.*, 1988), and a complete nucleotide sequence of *cfxF* from a prokaryote is not available. Only the sequences of the *cfxP* genes from spinach and *A. eutrophus* have been reported thus far (Kossmann *et al.*, 1990; Roesler & Ogren, 1988).

It thus appears that both PRK and FBPase have received only minor attention despite their important role in the Calvin cycle. We therefore set out to characterize the cloned PRK and FBPase structural genes from *Xanthobacter flavus* H4-14, a bacterium capable of autotrophic growth on methanol and molecular hydrogen (Lidstrom-O'Connor *et al.*, 1983; Meijer *et al.*, 1990*a*). In this paper we report the complete nucleotide sequences of these genes.

### Methods

**Bacterial strains and plasmids.** *Escherichia coli* JM101 (Yanisch-Perron *et al.*, 1985) was used as a host for pBLUESCRIPT (Vector Cloning Systems), M13mp18 and M13mp19 (Yanisch-Perron *et al.*, 1985) and their derivatives. pCD102 is a pVK100 cosmid, containing a 24 kb chromosomal DNA fragment from *X. flavus* H4-14, encoding FBPase and PRK (Lehmiche & Lidstrom, 1985; Meijer *et al.*, 1990*b*).

**Media and growth conditions.** *E. coli* strains were grown on LB medium at 37 °C (Maniatis *et al.*, 1982). Agar was added to 1.5% (w/v) to solidify the medium. When necessary the following supplements

---

† Present address: Department of Microbiology, The Ohio State University, 484 West 12th Avenue, Columbus, OH 43210-1292, USA.

**Abbreviations:** FBPase, fructosebisphosphatase; F-2,6-P, fructose 2,6-bisphosphate; LB, Luria broth; ORF, open reading frame; PRK, phosphoribulokinase; RuBP, ribulose bisphosphate; RuBisC/O, ribulose-1,5-bisphosphate carboxylase/oxygenase; RuMP, ribulose monophosphate.

The nucleotide sequence data reported in this paper have been submitted to EMBL and will appear in the EMBL/GenBank/DBJ Nucleotide Sequence Databases under the accession number X17252.




Fig. 1. Restriction map of the 2.5 kb DNA *Sal*I fragment from *X. flavus* H4-14 and the positions of *cfxF* and *cfxP*. Arrows indicate the direction of transcription. Restriction sites: Ba, *Bam*HI; Bg, *Bgl*II; Ec, *Eco*RI; Ps, *Pst*I; Sa, *Sal*I.

were added (in  $\mu\text{g ml}^{-1}$ ): ampicillin, 50; 5-bromo-4-chloro-3-indolyl  $\beta$ -D-galactoside, 20; tetracycline, 12.5. Isopropyl  $\beta$ -D-thiogalactoside was used at a concentration of 0.1 mM.

**DNA manipulations.** Plasmid DNA was isolated from *E. coli* by using either the Triton X-100 lysis method of Rodriguez & Tait (1983), in which the column chromatography step was omitted for preparative isolations, or the alkaline lysis method of Birnboim & Doly (1979) for isolations on a smaller scale. Restriction enzymes and T4 DNA ligase were used according to the manufacturer's instructions. Analysis of restriction digests by gel electrophoresis, recovery of DNA fragments from low-melting-point agarose and transformation of *E. coli* were carried out as described by Maniatis *et al.* (1982). Sequence determination was done as described by Sanger *et al.* (1977), except that 7-deaza-dGTP was used instead of dGTP, and Sequenase (United States Biochemical Corporation) was used instead of the Klenow fragment of DNA polymerase. When necessary, custom primers (Eurosequence, Groningen, the Netherlands) were used instead of the universal primer.

## Results and Discussion

### Nucleotide sequences of *cfxF* and *cfxP*

The localization of the *cfxF* and *cfxP* genes in *X. flavus* H4-14, and their cotranscription in the order *cfxFP*, has been established in previous work (Meijer *et al.*, 1990b; Fig. 1). A 2.5 kb *Sal*I fragment from pCD102 encoding FBPase and PRK was subcloned in pBLUESCIBE, yielding plasmid pWL5. This plasmid was digested with the appropriate restriction enzymes, and the restriction fragments were ligated into M13mp18 and M13mp19. The total nucleotide sequence was derived from both DNA strands and fully overlapped. The C + G content of the *Sal*I fragment was 66.4 mol%, which is close to the C + G content of the chromosomal DNA (68 mol%; Meijer *et al.*, 1990a).

Two open reading frames (ORFA, ORFB) were identified in the nucleotide sequence, each preceded by plausible ribosome-binding sites. Both were transcribed in the same direction as *cfxF* and *cfxP* (Meijer *et al.*, 1990b), and were separated by only 39 bp. ORFA had a length of 1092 bp, and would encode a protein of 38739 Da. ORFB had a length of 873 bp, encoding a

protein of 33409 Da. Due to the high C + G content, the codon usage in both ORFA and ORFB displayed a strong bias towards the use of codons having a C or G in the wobble position.

The deduced amino acid sequences derived from both ORFs were compared to 10856 entries in the SWPROT-database (Protein Database, Release 11, June 1989, University of Geneva, Switzerland), using the program FASTP (k-tuple = 2; Lipman & Pearson, 1985). On the basis of this comparison, ORFA and ORFB were identified as *cfxF* and *cfxP*, respectively. The deduced molecular masses of the *cfxF* and *cfxP* gene products corresponded well with the sizes of the *cfxF* gene product, expressed in *E. coli* (40 kDa; W. G. Meijer & L. Dijkhuizen, unpublished observations), and PRK from *X. flavus* H4-14 (33 kDa; Lehmicke & Lidstrom, 1985). The nucleotide sequences and deduced amino acid sequences of *cfxF* and *cfxP* are shown in Fig. 2.

### Comparison of FBPase primary structures

A comparison of the primary structures of FBPase from various sources is shown in Fig. 3. The FBPase amino acid sequence from *X. flavus* H4-14 displays 32 to 38% similarity with FBPases from heterotrophic sources, which show 41 to 63% similarity amongst each other (Table 1). Of the two autotrophic FBPase proteins, from wheat chloroplasts and *A. eutrophus*, only the latter shows a high similarity (52%) with the *X. flavus* H4-14 protein, although all three autotrophic FBPase proteins have the same physiological role. In contrast, the RuBisC/O proteins from plants are very similar to those from *A. eutrophus* and *X. flavus* H4-14 (Andersen & Caton, 1987; Meijer *et al.*, 1990b). An explanation for this difference in sequence conservation could be that the structural constraints in RuBisC/O, a hexadecamer consisting of two types of subunits, are higher than those in the FBPase protein, allowing less sequence variation. Alternatively, the FBPase proteins from eukaryotic and prokaryotic species may have evolved from different ancestors.

Recently the three-dimensional structure of pig kidney FBPase and its complexes with AMP and fructose 2,6-bisphosphate (F-2,6-P) have become available (Ke *et al.*, 1989). AMP and F-2,6-P inhibit FBPase activity in a number of organisms. AMP inhibition is allosteric, whereas F-2,6-P probably binds to the active site (Ke *et al.*, 1989; Liu *et al.*, 1989). The F-2,6-P-binding site consists of the following residues (numbering for pig kidney FBPase): Asn-212, Tyr-244, Gly-246, Ser-247, Met-248, Tyr-264 and Lys-274. These residues are either identical, or represented by a conservative replacement in the *X. flavus* H4-14 protein. Residues making up a negatively charged pocket, forming one wall of the

```

GGACCGGAAGACGACCAAAAAATAAACCTGTGACGGCCCGCTGAACGGGCCGCGGAGGAGCCCTCAGGGAGATCTGCCATGTTGGAGC
10 20 30 40 50 60 70 80 90
M L E P
N A D H R A A V A Q A A G V A A S R I T L T V M L D E W A G
CGAACGCAGACCATCGGGCCGAGTTGCCAGGCTGCAGGTGTGCGAGCCTCGCGTATCACGCTCACCGTCATGCTCGACGAGTGGGCGG
100 110 120 130 140 150 160 170 180
A D A R R R A V A D T V C A L A T G C A S L A A A I A E G P
GTGGGATGCCCGCCGCTGCCGTGCGGATACCGTCTGCGCCCTCGCCACCGGTTGCGCGTCGCTGGCCGACGCCATCGCCGAGGGGC
190 200 210 220 230 240 250 260 270
L A G D L A R T L S S G E A G E G Q K A L D V I S N D I V I
CGCTCGCGCGATCTCGCCGACCCCTCTCCCTCGGCGAGGCGGCGAAGGCCGAGGCGTGGACGTCATCTCCAACGACATCGTCA
280 290 300 310 320 330 340 350 360
G A L K A A P V A A V A S E E N D A P V L L D P T A P L L V
TCGGCGCGCTGAAGGCGGCGCGGTGCGGCGGTGCGATCCGAGGAGAAGACGCGCCCGGCTCGTCTGATCCACCGCGCGCTGCTCG
370 380 390 400 410 420 430 440 450
A I D P L D G S S N I D T D I S V G T I F A V F P R P E G A
TCGCCATCGATCCGCTGGACGCGCTCGTCAACATCGACACCGACATCTCGGTGCGCACCATCTCGCGCTCTCCGCGTCCCGAGGGG
460 470 480 490 500 510 520 530 540
D A S E P S A F L Q N G R D M L A A G Y V I Y G P H T A M M
CCGACGCTCCGAACCCCTCTGCCCTTCTGCAGAACGGCGGGACATGCTGGCCGCGGTTATGTGATCTACGGCCCCACACCGCATGA
550 560 570 580 590 600 610 620 630
L T L G A G T W H F A L D R A G L F R L V D A E V K V K E G
TGCTGACGCTCGGCGCGCACCTGGCACTTCGCGCTCGACCGCGCGGCTCTTCGCGCTGGTCGATGACGAGGTGAAGTGAAGGAGG
640 650 660 670 680 690 700 710 720
A A E F A I N M S N Y H H W D V P V R D Y V D D C L A G K K
GCGCGCGGATTCGCGCATCAACATGTCGAACCTACCACTGGGACGTGCGGTCGCTGACTATGTGGATGACTGCGCTCGCGGCAAGA
730 740 750 760 770 780 790 800 810
G P R E R D F N M R W V A S M V A D A H R I F Q R G G I Y L
AGGGCCCGCGGAGCGCGACTTCAACATGGCGTGGTGGCGCTCCATGGTGGCGGACGCGCATCGCATCTTCCAGCGCGCGGCGCATATC
820 830 840 850 860 870 880 890 900
Y P G D G R K G Y T H G R L R L L Y E A F P V A F L M E Q A
TCTATCCGGCGATGGCGGAAGGGCTACACGCAACGGCGCGCTGCGCTCTCTACGAGGCTTCCCGCTGCGCTTCTGATGGAGCAGG
910 920 930 940 950 960 970 980 990
S G S A T D G R G A I L D L S A T G L H Q R V P F I F G S R
CGAGCGGTTCGCGCAACCGCGCGGCGGCACTTCTGGACCTTTCGCGCACCGGCTGACCGAGCGGTGCGGTTCTCTCGGCTCCC
1000 1010 1020 1030 1040 1050 1060 1070 1080
D E V A R V S R Y H L E P N G H G E R S P L F A R R G L F I
GCGACGAGGTGGCGCGGCTCTCCCGTATACCTGGAGCGCAACGGCCATGGCGAGCGCTCGCGCTGTTTCGCGCGCGCGGACTGTTCA
1090 1100 1110 1120 1130 1140 1150 1160 1170
M S I K H P I I V V T G S S G A G
TCTGAACCGCGCCCCCTGATCTCGGAGATTGAGCACACCATGTCCATCAAGCACCCCATCATGTGCTGACCGGTTCTCGGGCGCGGG
1180 1190 1200 1210 1220 1230 1240 1250 1260
T T S V K R T F E Q I F Y R E K V K A A F V E G D S F H R Y
AACGACCTCGGTGAGCGGACCTTCGAGCATCTTCTATCGCGAGAGGTCAAGGCGGCTTCGTGGAAGCGACAGCTTCCACCGCTA
1270 1280 1290 1300 1310 1320 1330 1340 1350
D R Y E M R E L M A A E A A K G N K H F S H F S P E T N R L
CGACGCTGAGATGCGGAGCTGATGGCGCGGCGGCAAGGCGCAACGACCTTCAGCCACTTCTGCGCGAGACCAACCGGCT
1360 1370 1380 1390 1400 1410 1420 1430 1440
D D L A Q L F K D Y G A T G S G R F R H Y V H D A G E A K L
CGACGATCTGGCCGAGCTGTTCAAGGACTATGGGGGACCGGCTCCGGCGCTTCGGGCACTATGTCACGATGCCGCGAAGCGAAGCT
1450 1460 1470 1480 1490 1500 1510 1520 1530
Y N T E P G R F T D W E D L E Q G T D I L F Y E G L H G A V
GTACATACCGAGCCCGCGCTTACCGACTGGGAAGATCTGGAGCAGGCGACCGACATCTCTTACGAGGGGCTGCACGGGCGGT
1540 1550 1560 1570 1580 1590 1600 1610 1620
V T D E L N L A Q H A D L K I G V V P V I N L E W I Q K I H
CGTCACCGAAGCTGAACCTCGCCGAGCATGCCGACCTGAAGATCGGCGTGGTGGCGGTGATCAACCTGGAGTGGATCCAGAAGATCCA
1630 1640 1650 1660 1670 1680 1690 1700 1710
R D K A T R G Y T T E D V T D T I M R R M P D Y V R Y I C P
TCGGACAGGCGCACCGGCTACACCGAGGACGTGACCGACCATCATGCGGCGCATGCCGATTACGTGCGCTACATCTGCCC
1720 1730 1740 1750 1760 1770 1780 1790 1800
Q F T E T D I N F Q R V P T V D T S N P F V A R W I P T P D
GCAGTTACCGGAGACCGACATCAACTTCCAGCGGTGCCGAGGTGGACACCTCCAACCGGTTCTGTCGCGCGCTGGATCCCGAGCGCGGA
1810 1820 1830 1840 1850 1860 1870 1880 1890
E S M V V I R F R D P H G I D F P Y L L S M I H N S F M S R
CGAATCGATGTCGTGATCCGCTTCCGCGACCGCGACGGCATGATTTCCCTATTGCTGTCGATGATCCACAACAGCTTCATGTCGCG
1900 1910 1920 1930 1940 1950 1960 1970 1980
A N S I V I P G N K Q D L A M Q L L L T P L I M K L M D R K
GGCGAATTCCATCGTCATCCCGGGCAACAAGCAGGATCTCGCCATGCAAGCTCTTCTGACCCCGCTCATCATGAAGCTGATGGACAGGAA
1990 2000 2010 2020 2030 2040 2050 2060 2070
R R A G
GCGCGAGCCGCGTGAAGCGGCGCGAAGCTCCC
2080 2090 2100

```

Fig. 2. Nucleotide sequence and deduced amino acid sequence of *cfxF* and *cfxP* from *X. flavus* H4-14. The first open reading frame represents *cfxF* and the second *cfxP*. The amino acids are represented by the one-letter code. The putative ribosome-binding sites are underlined.




Fig. 3. Alignment of FBPase proteins from: X, *X. flavus* H4-14; A, *Alcaligenes eutrophus* (Kossmann *et al.*, 1990); E, *Escherichia coli* (Hamilton *et al.*, 1988); S, *Schizosaccharomyces pombe* (Rogers *et al.*, 1988); Y, *Saccharomyces cerevisiae* (Rogers *et al.*, 1988); W, wheat chloroplast (Raines *et al.*, 1988); P, pig kidney (Marcus *et al.*, 1982). The mature form of the chloroplast FBPase was used in the alignment. ■, identical residue; ., conservative substitution following the scheme: PAGST, QNED, ILMV, HKR, YFW, C. Only similarities between more than two sequences are indicated. Numbers in parentheses refer to amino acids. Except for the terminal residues, only the residues that differ from the *X. flavus* H4-14 sequence are indicated.

Table 1. Similarity between FBPase primary structures

Similarities are in percentages. X, *Xanthobacter flavus* H4-14; A, carboxy terminus from *Alcaligenes eutrophus* (Kossmann *et al.*, 1990); E, *Escherichia coli* (Hamilton *et al.*, 1988); W, wheat chloroplast (Raines *et al.*, 1988); P, pig kidney (Marcus *et al.*, 1982); Y, *Saccharomyces cerevisiae* (Rogers *et al.*, 1988); S, *Schizosaccharomyces pombe* (Rogers *et al.*, 1988).

	X	A	E	W	P	Y	S
X	100						
A	52	100					
E	38	38	100				
W	34	39	46	100			
P	36	36	45	46	100		
Y	35	36	43	41	47	100	
S	32	40	46	43	46	63	100

F-2,6-P-binding site, Asp-118, Asp-121, Glu-280, Glu-97, Glu-98, Lys-71 and Arg-276 (numbering for pig kidney FBPase) are identical in the *X. flavus* H4-14 FBPase sequence.  $Mg^{2+}$ , which is essential for FBPase activity, interacts with both the negatively charged pocket and a phosphate group from F-2,6-P. This

strongly suggests that F-2,6-P binds to the active site (Ke *et al.*, 1989). Residues close to the AMP-binding site in the pig FBPase sequence are either not conserved or are represented by a conservative substitution in the *X. flavus* H4-14 sequence.

The fact that the F-2,6-P-binding site residues are strongly conserved in the *X. flavus* H4-14 FBPase sequence indicates that this region is indeed essential for FBPase activity, and thus may constitute the active site. Residues close to the AMP-binding site are not conserved. The *Nocardia opaca* FBPase, which is specifically induced during autotrophic growth, is not very sensitive to AMP. In contrast, the FBPase isoenzyme functioning in gluconeogenesis in the same organism is fully inhibited by AMP at concentrations of 100  $\mu M$  (Amachi & Bowien, 1979). The properties of the enzyme in *X. flavus* H4-14 are currently under investigation.

Characteristics which are specific for a particular FBPase, such as the insertions in the *Schizosaccharomyces pombe* and wheat chloroplast FBPase sequences,




Fig. 4. Alignment of PRK proteins from: X, *X. flavus* H4-14; A, *Alcaligenes eutrophus* (Kossmann *et al.*, 1990); R, *Rhodobacter sphaeroides* (Hallenbeck & Kaplan, 1987); S, spinach (Roesler & Ogren, 1988). ■, identical residue; ., conservative substitution as in Fig. 3. Only similarities between more than two sequences are indicated. Regions of identity are underlined (see text). Numbers in parentheses refer to amino acids. Except for the terminal residues, only the residues that differ from the *X. flavus* H4-14 sequence are indicated.

and the cAMP-dependent protein kinase consensus sequence in the *Saccharomyces cerevisiae* amino terminus, are not conserved in the *X. flavus* H4-14 sequence (Fig. 3; Raines *et al.*, 1988; Rogers *et al.*, 1988).

#### Comparison of PRK primary structures

As was observed for the FBPase protein, the PRK amino acid sequence from *X. flavus* H4-14 is not very similar (less than 22% identity) to that from a eukaryotic plant, in this case spinach (Fig. 4). In contrast, the *X. flavus* H4-14 and *A. eutrophus* proteins display 65% similarity. This does not come as a surprise, as the quaternary structure of prokaryotic and eukaryotic PRK proteins is very different. In bacteria, PRK is present as an octamer of 33 kDa subunits, whereas in spinach, PRK is a dimer of 45 kDa subunits (Krieger & Miziorko, 1986; Rippel & Bowien, 1984; Siebert & Bowien, 1984). This indicates that residues important in subunit interaction in the prokaryotic holoenzyme are not conserved in the eukaryotic proteins.

In the amino-terminal protein sequence of PRK of both prokaryotic and eukaryotic origin a sequence can be recognized which is supposed to be an ATP-binding site (region A, Fig. 4; Hallenbeck & Kaplan, 1987; Klintonworth *et al.*, 1985; Porter *et al.*, 1988). The spinach PRK sequence contains two unique cysteine residues, one at position 16, within the proposed ATP-binding site consensus sequence, the other at position 55. These cysteine residues can be oxidized to form a disulphide bridge, causing inactivation of PRK (Porter *et al.*, 1988). The oxidation and alkylation of Cys-16 by (bromoacetyl) ethanolamine phosphate can be prevented by ATP, strengthening the hypothesis that Cys-16 is at the active

site (Porter & Hartman, 1986). The functions of the other conserved regions (B and C: Fig. 4) remain to be established, although they may play a role in the binding of the substrate, RuMP. The identified conserved regions may be suitable targets for mutagenesis to elucidate their function.

The authors thank Dr B. Bowien for making results available prior to publication, and Professor Dr W. Harder for valuable discussions. The investigations were supported by the Foundation for Fundamental Biological Research (BION), which is subsidized by the Netherlands Scientific Organization (NWO).

#### References

- AMACHI, T. & BOWIEN, B. (1979). Characterization of two fructose bisphosphatase isoenzymes from the hydrogen bacterium *Nocardia opaca* 1b. *Journal of General Microbiology* **113**, 347–356.
- ANDERSEN, K. & CATON, J. (1987). Sequence analysis of the *Alcaligenes eutrophus* chromosomally encoded ribulose bisphosphate carboxylase large and small subunit and their gene products. *Journal of Bacteriology* **169**, 4547–4558.
- BIRNBOIM, H. C. & DOLY, J. (1979). A rapid alkaline extraction procedure for screening of recombinant plasmid DNA. *Nucleic Acids Research* **7**, 1513–1523.
- GIBSON, J. L. & TABITA, F. R. (1987). Organization of phosphoribulokinase and ribulose bisphosphate carboxylase/oxygenase genes in *Rhodospseudomonas (Rhodobacter) sphaeroides*. *Journal of Bacteriology* **169**, 3685–3690.
- GIBSON, J. L. & TABITA, F. R. (1988). Localization and mapping of CO<sub>2</sub> fixation genes within two gene clusters in *Rhodobacter sphaeroides*. *Journal of Bacteriology* **170**, 2153–2158.
- HALLENBECK, P. L. & KAPLAN, S. (1987). Cloning of the gene for phosphoribulokinase activity from *Rhodobacter sphaeroides* and its expression in *Escherichia coli*. *Journal of Bacteriology* **169**, 3669–3678.
- HAMILTON, W. D. O., HARRISON, D. A. & DYER, T. A. (1988). Sequence of the *Escherichia coli* fructose-1,6-bisphosphatase gene. *Nucleic Acids Research* **16**, 8707.

- KE, H., THORPE, C. H., SEATON, B. A., MARCUS, F. & LIPSCOMB, W. N. (1989). Molecular structure of fructose-1,6-bisphosphate at 2.8-Å resolution. *Proceedings of the National Academy of Sciences of the United States of America* **86**, 1475–1479.
- KLINTWORTH, R., HUSEMANN, M., SALNIKOW, J. & BOWIEN, B. (1985). Chromosomal and plasmid locations for phosphoribulokinase genes in *Alcaligenes eutrophus*. *Journal of Bacteriology* **164**, 954–956.
- KLINTWORTH, R., HUSEMANN, M., WEISSENBORN, C. & BOWIEN, B. (1988). Expression of the plasmid-encoded phosphoribulokinase gene from *Alcaligenes eutrophus*. *FEMS Microbiology Letters* **49**, 1–6.
- KOSSMANN, J., KLINTWORTH, R. & BOWIEN, B. (1990). Sequence analysis of the chromosomal and plasmid genes encoding phosphoribulokinase from *Alcaligenes eutrophus*. *Gene* **85**, 247–252.
- KRIEGER, T. J. & MIZIORKO, H. M. (1986). Affinity labelling and purification of spinach leaf ribulose-5-phosphate kinase. *Biochemistry* **25**, 3496–3501.
- LEHMICKE, L. G. & LIDSTROM, M. E. (1985). Organization of the genes necessary for growth of the hydrogen-methanol autotroph *Xanthobacter* sp. strain H4-14 on hydrogen and carbon dioxide. *Journal of Bacteriology* **162**, 1244–1249.
- LIDSTROM-O'CONNOR, M. E., FULTON, G. L. & WOPAT, A. E. (1983). *Methylobacterium ethanolicum*: a syntrophic association of two methylotrophic bacteria. *Journal of General Microbiology* **129**, 3139–3148.
- LIPMAN, D. J. & PEARSON, W. R. (1985). Rapid and sensitive protein similarity searches. *Science* **227**, 1435–1441.
- LIU, F., ROY, M. & FROMM, H. J. (1989). The site of substrate and fructose 2,6-bisphosphate binding to rabbit liver fructose-1,6-bisphosphatase. *Biochemical and Biophysical Research Communications* **161**, 689–695.
- MANIATIS, T., FRITSCH, E. F. & SAMBROOK, J. (1982). *Molecular Cloning: a Laboratory Manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- MARCUS, F., EDELSTEIN, I., REARDON, I. & HEINRIKSON, R. L. (1982). Complete amino acid sequence of pig kidney fructose-1,6-bisphosphatase. *Proceedings of the National Academy of Sciences of the United States of America* **79**, 7161–7165.
- MEIJER, W. G., CROES, L. M., JENNI, B., LEHMICKE, L. G., LIDSTROM, M. E. & DIJKHUIZEN, L. (1990a). Characterization of *Xanthobacter* strains H4-14 and 25a and enzyme profiles after growth under autotrophic and heterotrophic conditions. *Archives of Microbiology* **153**, 360–367.
- MEIJER, W. G., ARNBERG, A. C., ENEQUIST, H. G., TERPSTRA, P., LIDSTROM, M. E. & DIJKHUIZEN, L. (1990b). Identification and organization of carbon dioxide fixation genes in *Xanthobacter flavus* H4-14. *Molecular and General Genetics* (in the Press).
- PORTER, M. A. & HARTMAN, F. C. (1986). Commonality of catalytic and regulatory sites of spinach phosphoribulokinase: characterization of a tryptic peptide that contains an essential cysteinyl residue. *Biochemistry* **25**, 7314–7318.
- PORTER, M. A., STRINGER, C. D. & HARTMAN, F. C. (1988). Characterization of the regulatory thioredoxin site of phosphoribulokinase. *Journal of Biological Chemistry* **263**, 123–129.
- RAINES, C. A., LLOYD, J. C., LONGSTAFF, M., BRADLEY, D. & DYER, T. A. (1988). Chloroplast fructose-1,6-bisphosphatase: the product of a mosaic gene. *Nucleic Acids Research* **16**, 7931–7942.
- RIPPEL, S. & BOWIEN, B. (1984). Phosphoribulokinase from *Rhodospseudomonas acidophila*. *Archives of Microbiology* **139**, 207–212.
- RODRIGUEZ, R. L. & TAIT, R. C. (1983). *Recombinant DNA Techniques: an Introduction*. Reading, Massachusetts: Addison-Wesley.
- ROESLER, K. R. & OGREN, W. L. (1988). Nucleotide sequence of spinach cDNA encoding phosphoribulokinase. *Nucleic Acids Research* **16**, 7192.
- ROGERS, D. T., HILLER, E., MITSOCK, L. & ORR, E. (1988). Characterization of the gene for fructose-1,6-bisphosphatase from *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*. *Journal of Biological Chemistry* **263**, 6051–6057.
- SANGER, F., NICKLEN, S. & COULSON, A. R. (1977). DNA sequencing with chain terminating inhibitors. *Proceedings of the National Academy of Sciences of the United States of America* **74**, 5463–5467.
- SIEBERT, K. & BOWIEN, B. (1984). Evidence for an octameric structure of phosphoribulokinase from *Alcaligenes eutrophus*. *Biochimica et Biophysica Acta* **787**, 208–214.
- TABITA, F. R. (1988). Molecular and cellular regulation of autotrophic carbon dioxide fixation in microorganisms. *Microbiological Reviews* **52**, 155–189.
- YANISCH-PERRON, C., VIEIRA, J. & MESSING, J. (1985). Improved M13 phage cloning vectors and host strains. Nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**, 103–119.